

## Combined pill (oestrogen and progestogen) with a break : what to do when missing a pill?

Missing more than 12 hours, 1 pill of your packet

1 <sup>rst</sup> week	2 <sup>nd</sup> week	3 <sup>rd</sup> week	4 <sup>th</sup> week
000000	000000	000000	
• take the missed pill as soon as possible + take the pill of the day at the usual time	• take the missed pill as soon as possible + take the pill of the day at the usual time	• take the missed pill as soon as possible + take the pill of the day at the usual time	7 days break
<ul><li>keep taking the following pills</li></ul>	• keep taking the following pills	• keep taking the following pills	
• use condom for 7 days			
If you have had sexual intercourse within the last 5 days:	• the use of <b>condoms for 7 days</b> avoids the need for emergency contraception in the event of a 2 <sup>nd</sup> omission	• the use of <b>condoms for 7 days</b> avoids the need for emergency contraception in the event of a 2nd omission	
• take an emergency contraception as fast as you can		• start a new packet of pills straightaway,	
Norlevo® (1 <sup>rs</sup> choice): use condom for 7 days and keep on with your regular contraceptive pill		with no pill-free days	
ellaOne®: if possible, stop your contraceptive pill for 5 days, then start a new package + use condom during all the tablet			

Missing 2 pills or more:

Go immediately to a sexual health center, pharmacy, maternity hospital or to your gynecologist